



E-ISSN: 2709-9407  
 P-ISSN: 2709-9393  
 JMPES 2020; 1(1): 10-19  
 © 2020 JMPES  
[www.mathematicaljournal.com](http://www.mathematicaljournal.com)  
 Received: 05-11-2019  
 Accepted: 10-12-2019

**Tulkin Mamatov**  
 Department of Higher  
 mathematics, Bukhara  
 Technological Institute of  
 Engineering, Bukhara,  
 Uzbekistan

**Nemat Mustafoev**  
 Department of Higher  
 mathematics, Bukhara  
 Technological Institute of  
 Engineering, Bukhara,  
 Uzbekistan

## Estimates of the zygmond type of mixed fractional integrals of riemann-liouville and derivatives of marchaud

**Tulkin Mamatov and Nemat Mustafoev**

### Abstract

Zygmund type estimates are obtained for the mixed continuity modulus of some mixed fractional integrals. It is known, the Riemann-Liouville fractional integration operator establishes an isomorphism between Hölder spaces for functions of one variable. We study mixed Riemann-Liouville fractional integration operators and mixed fractional derivative in Marchaud form of function of two variables.

**Keywords:** functions of two variables, fractional derivative of Marchaud form, mixed fractional derivative, mixed fractional integral, mixed continuity modulus

### 1. Introduction

One of the most important problems in the theory of integral operators in space is the problem of elucidating the dependence of the smoothness of the image on the smoothness of the preimage. The solution to such a problem plays an important role in the solvability of integral equations, their stability and so on. The concept of smoothness can be formulated in a variety of terms. One of the ways of sufficiently fine-graining the smoothness of functions is the notion of generalized Hölderiness, formulated in terms of the behavior of the modulus of continuity. Thus one of the important questions in the theory of operators is as follows:

Let be  $A$  an operator acting in a Banach spaces  $X$  and let be the modulus of continuity  $\omega(f, h) = \sup_{|t| \leq h} \|f(x+h) - f(x)\|_X$  of  $X$ . How can the behavior of the modulus of

continuity be characterized  $\omega(A\varphi, h)$  if the behavior of the modulus of continuity of function  $\omega(\varphi; h): \omega(\varphi; h) \leq C\psi(h)$  for all is known  $\varphi \in X$ , where is  $\psi(x)$  a given continuous function,  $\psi(0) = 0$ . A similar problem can be considered completely solved for different spaces and also for the Hölder spaces of functions of one variable and weights, when fractional integration and fractional differentiation operators [1]. The assertion for multidimensional case for a mixed fractional Riemann-Liouville integral was studied [2, 15]. When mixed fractional derivatives form Marchaud

$$\begin{aligned} (D_{a+,c+}^{\alpha,\beta} \varphi)(x, y) &= \frac{(x-a)^{-\alpha} (y-c)^{-\beta}}{\Gamma(1-\alpha)\Gamma(1-\beta)} \varphi(x, y) + \\ &+ \frac{\alpha\beta}{\Gamma(1-\alpha)\Gamma(1-\beta)} \int_a^x \int_c^y \frac{\varphi(x, y) - \varphi(t, s)}{(x-t)^{1+\alpha} (y-s)^{1+\beta}} dt ds, \quad x > a, y > c \end{aligned} \quad (1)$$

were not studied. This paper is devoted to the study of the properties for functions of two variables. Consider the operator (1) in a rectangle  $Q = \{(x, y): a < x < b, c < y < d\}$ .

### 2. Preliminary

**Definition 1.** Let given bounded on  $[a, b]$  function  $\varphi(x)$ . Under modulus of continuity  $\omega(\varphi)$  understood expression

$$\sup_{h \in [0, \delta]} |\varphi(x+h) - \varphi(x)| = \omega(\varphi; \delta), \quad 0 < \delta \leq b-a.$$

1)

**Correspondence**  
**Tulkin Mamatov**  
 Department of Higher  
 mathematics, Bukhara  
 Technological Institute of  
 Engineering, Bukhara,  
 Uzbekistan

**Definition 2.** We denote by  $\Phi^1$  function class  $\omega(\delta) \in (0, b-a]$  and satisfying the conditions

- 1)  $\omega(\delta) > 0$  in  $(0, b-a]$ ,  $\lim_{\delta \rightarrow 0} \omega(\delta) = 0$ ;
- 2)  $\omega(\delta) \uparrow$  in  $(0, b-a]$ ;
- 3)  $\omega(\delta_1 + \delta_2) \leq \omega(\delta_1) + \omega(\delta_2)$ .

Below in the estimates we need inequalities:

1) if  $\omega(\varphi; h)$  is modulus continuity, then we have :  $x_2 \omega(\varphi; x_1) \leq C x_1 \omega(\varphi; x_2)$ ,  $x_2 \leq x_1$ ; (2)

2) if  $\lambda \leq 1$  then  $|x_1^\lambda - x_2^\lambda| \leq C(x_1 - x_2)x_2^{\lambda-1}$ ,  $x_1 \geq x_2 > 0$ ; (3)

3) if  $\lambda \geq 0$  then  $|x_1^\lambda - x_2^\lambda| \leq C(x_1 - x_2)x_1^{\lambda-1}$ ,  $x_1 \geq x_2 > 0$ . (4)

For a continuous function  $\varphi(x, y)$  on  $\mathbb{R}^2$  we introduce the notation

$$\left( \Delta_h \varphi \right)^{(1,0)}(x, y) = \varphi(x + h, y) - \varphi(x, y), \quad \left( \Delta_\eta \varphi \right)^{(0,1)}(x, y) = \varphi(x, y + \eta) - \varphi(x, y),$$

$$\left( \Delta_{h,\eta} \varphi \right)^{(1,1)}(x, y) = \varphi(x + h, y + \eta) - \varphi(x + h, y) - \varphi(x, y + \eta) + \varphi(x, y),$$

so that

$$\varphi(x + h, y + \eta) = \left( \Delta_{h,\eta} \varphi \right)^{(1,1)}(x, y) + \left( \Delta_h \varphi \right)^{(1,0)}(x, y) + \left( \Delta_\eta \varphi \right)^{(0,1)}(x, y) + \varphi(x, y). \tag{5}$$

Everywhere in the sequel by  $C, C_1, C_2$  etc we denote positive constants which may different values in different occurrences and even in the same line.

Now we introduce the following characteristics:

1) Private modules of continuity

$$\omega(\varphi; \delta, 0) = \sup_y \sup_{0 \leq h \leq \delta} \left| \left( \Delta_h \varphi \right)^{(1,0)}(x, y) \right| \quad \text{and} \quad \omega(\varphi; 0, \sigma) = \sup_x \sup_{0 \leq \eta \leq \sigma} \left| \left( \Delta_\eta \varphi \right)^{(0,1)}(x, y) \right|;$$

2) Mixed modulus continuity of order 1.1

$$\omega(\varphi; \delta, \sigma) = \sup_{x,y} \sup_{\substack{0 \leq h \leq \delta \\ 0 \leq \eta \leq \sigma}} \left| \left( \Delta_{h,\eta} \varphi \right)^{(1,1)}(x, y) \right|, \quad \text{where } 0 < \delta \leq b, 0 < \sigma \leq d.$$

It follows from the definition  $\omega(\varphi; \delta, \sigma)$  that this function belongs in each variable  $\Phi^1$ . In addition, we note that there is an inequality

$$\omega(\varphi; \delta, \sigma) \leq 2 \min \left\{ \omega(\varphi; \delta, 0), \omega(\varphi; 0, \sigma) \right\}. \tag{6}$$

**Definition 3.** We denote by  $\Phi^{1,1}$  the class of functions of two variables  $\omega(\delta, \sigma)$  satisfying conditions:

- 1)  $\omega(\delta, \sigma)$  in  $\delta$  for any fixed  $\sigma$ ;
- 2)  $\omega(\delta, \sigma)$  in  $\sigma$  for any fixed  $\delta$ .

We call this class the class of mixed modulus of continuity of the first order of continuity functions of two variables.

The following statements are known (see [1, p. 249-253]). We use the schemes of the proofs to make the presentation easier for two-dimensional case.

The following two theorems give estimates which might be called Zygmund types by analogy with the Zygmund estimate known in the theory of singular integrals and estimating the continuity modulus  $\omega(H\varphi, h)$  of a conjugate function  $H\varphi$  via the continuity modulus  $\omega(\varphi, h)$  of a function  $\varphi(x)$  itself.

Consider the one-dimensional fractional Riemann-Liouville integral

$$\left( I_{a+}^\alpha \varphi \right)(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} \varphi(t) dt, \quad x > a, 0 < \alpha < 1 \tag{7}$$

**Theorem 1.** [1]. Let  $\varphi(x)$  be continuous on  $[a, b]$  and let  $\varphi(a) = 0$ . For a fractional integral  $I_{a+}^\alpha \varphi$ ,  $0 < \alpha < 1$ , the estimate

$$\omega(I_{a+}^\alpha \varphi, h) \leq Ch \int_h^{b-a} \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt \tag{8}$$

is valid.

**Proof.** Representing (7) as

$$(I_{a+}^\alpha \varphi)(x) = \frac{\varphi(a)}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} dt + \frac{1}{\Gamma(\alpha)} \int_a^x \frac{\varphi(t) - \varphi(a)}{(x-t)^{1-\alpha}} dt = \Delta_1(x) + \Delta_2(x)$$

Let  $h > 0; x, x+h \in [a, b]$ . We have

$$\begin{aligned} \Delta_2(x+h) - \Delta_2(x) &= \frac{\varphi(x) - \varphi(a)}{\Gamma(1+\alpha)} [(x+h-a)^\alpha - (x-a)^\alpha] + \frac{1}{\Gamma(\alpha)} \int_0^h \frac{\varphi(x+t) - \varphi(t)}{(h-t)^{1-\alpha}} dt + \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^{x-a} [\varphi(x-t) - \varphi(t)] [(h+t)^{\alpha-1} - t^{\alpha-1}] dt = I_1 + I_2 + I_3 \end{aligned}$$

We have:  $|I_1| \leq C\omega(\varphi, x) |(x+h-a)^\alpha - (x-a)^\alpha|$ . In the case  $x-a \leq h$  we have  $|I_1| \leq Ch^\alpha \omega(\varphi, h)$ . Let  $x-a \geq h$  then

$$|I_1| \leq C\omega(\varphi, x-a) (x-a)^\alpha \left[ \left(1 + \frac{h}{x-a}\right)^\alpha - 1 \right] \leq Ch(x-a)^{\alpha-1} \omega(\varphi, x-a) \tag{9}$$

Since

$$C(x-a)^{\alpha-1} \omega(\varphi, x-a) \leq \omega(\varphi, x-a) \int_{x-a}^{b-a} t^{\alpha-2} dt \leq \int_{x-a}^{b-a} \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt \leq \int_h^{b-a} \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt$$

$$|I_1| \leq Ch \int_h^{b-a} \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt$$

It follows from (9) that

$$|I_2| \leq \int_0^h \frac{\omega(\varphi, t) dt}{(h-t)^{1-\alpha}} = h^\alpha \int_0^1 \frac{\omega(\varphi, h\xi)}{(1-\xi)^{1-\alpha}} d\xi \leq Ch^\alpha \omega(\varphi, h) \quad C = \int_0^1 (1-\xi)^{\alpha-1} d\xi$$

Further with

To estimate  $I_3$  we distinguish the case 1)  $x-a \geq h$  and 2)  $x-a \leq h$ . In the first case

$$|I_3| \leq C \left\{ \int_0^h \omega(\varphi, t) [t^{\alpha-1} - (h+t)^{\alpha-1}] dt + \int_h^{x-a} \omega(\varphi, t) [t^{\alpha-1} - (h+t)^{\alpha-1}] dt \right\} \leq C_2 \left[ h^\alpha \omega(\varphi, h) + h \int_h^{b-a} \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt \right]$$

Obviously in the second case  $|I_3| \leq C_1 h^\alpha \omega(\varphi, h)$ .

Estimates for  $I_1, I_2, I_3$  lead to (8) if we take into account the fact that  $h^\alpha \omega(\varphi, h)$  is dominated by the right-hand side of (8).

The latter is easily obtained in view of the monotonicity of the function  $\omega(\varphi, t)$ .

The Marchaud fractional differentiation operator has a form

$$(D_{a+}^\alpha \varphi)(x) = \frac{\varphi(x)}{(x-a)^\alpha \Gamma(1-\alpha)} + \frac{\alpha}{\Gamma(1-\alpha)} \int_a^x \frac{\varphi(x) - \varphi(t)}{(x-t)^{1+\alpha}} dt \tag{10}$$

where  $0 < \alpha < 1$ .

**Theorem 2.** Let  $\varphi(x)$  be continuous on  $[a, b]$  and  $\varphi(a) = 0$ . Then its fractional derivative  $D_{a+}^\alpha \varphi, 0 < \alpha < 1$  admits the estimate

$$\omega(D_{a+}^\alpha \varphi, h) \leq C \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt \tag{11}$$

provided that the integral on the right-hand side converges.

**Proof.** We present (10) as

$$\begin{aligned} (D_{a+}^\alpha \varphi)(x) &= \frac{\varphi(a)}{(x-a)^\alpha \Gamma(1-\alpha)} + \frac{\varphi(x) - \varphi(a)}{(x-a)^\alpha \Gamma(1-\alpha)} + \frac{\alpha}{\Gamma(1-\alpha)} \int_a^x \frac{\varphi(x) - \varphi(t)}{(x-t)^{1+\alpha}} dt = \\ &= \frac{\varphi(a)}{\Gamma(1-\alpha)} (x-a)^{-\alpha} + F_1(x) + F_2(x) \end{aligned}$$

$$F_1(x) = \frac{\varphi(x) - \varphi(a)}{(x-a)^\alpha}, \quad 0 < \alpha < 1$$

We begin by noting that the function  $F_1(x)$  admits the estimate

$$\omega(F_1, h) \leq C \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt \tag{12}$$

Let us prove (12). Taking  $h > 0$  we have

$$F_1(x+h) - F_1(x) = [\varphi(x) - \varphi(a)] \left[ (x+h-a)^{-\alpha} - (x-a)^{-\alpha} \right] + \frac{\varphi(x+h) - \varphi(x)}{(x+h-a)^\alpha} = A_1 + A_2$$

Hence

$$|A_2| \leq (x+h-a)^{-\alpha} \omega(\varphi, h) \leq h^{-\alpha} \omega(\varphi, h) \leq C \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt$$

here in the last inequality we made use of the fact that the function  $\frac{\omega(\varphi, t)}{t}$  almost decreases. Further again taking this decreasing into account for  $A_1$  when  $x-a \leq h$  we have

$$|A_1| \leq (x-a)^{-\alpha} \omega(\varphi, x-a) \leq C \int_0^{x-a} t^{-1-\alpha} \omega(\varphi, t) dt \leq C \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt$$

When  $x-a \geq h$  the mean value theorem yields the estimate

$$|A_1| \leq Ch(x-a)^{-1-\alpha} \omega(\varphi, x-a) \leq C \frac{\omega(\varphi, h)}{h^\alpha} \leq C \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt$$

Gathering estimates for  $A_1$  and  $A_2$  we obtain the inequality (12).

To prove the theorem it's sufficient in view of (12) to consider only the second summand in the expression (10) for the Marchaud fractional derivative. For this function we have

$$\begin{aligned} F_2(x+h) - F_2(x) &= \int_a^x \frac{\varphi(x+h) - \varphi(x)}{(x+h-t)^{1+\alpha}} dt + \int_{x-a}^{x+h-a} \frac{\varphi(x+h) - \varphi(t)}{(x+h-t)^{1+\alpha}} dt + \\ &+ \int_a^x (f(x) - f(t)) \left[ (x+h-t)^{-1-\alpha} - (x-t)^{-1-\alpha} \right] dt = J_1 + J_2 + J_3 \end{aligned}$$

If  $x-a \leq h$  then  $|J_1| = \int_0^{x-a} \frac{\varphi(x+h) - \varphi(x)}{(h+t)^{1+\alpha}} dt \leq \int_0^{x-a} \frac{\omega(\varphi, h)}{(h+t)^{1+\alpha}} dt \leq \int_0^h \frac{\omega(\varphi, h)}{(h+t)^{1+\alpha}} dt \leq C \frac{\omega(\varphi, h)}{h^\alpha}$

If  $x-a \geq h$  we have

$$|J_1| \leq \int_0^{x-a} \frac{\omega(\varphi, h)}{(h+t)^{1+\alpha}} dt \leq \int_0^h \frac{\omega(\varphi, h)}{(h+t)^{1+\alpha}} dt + \int_h^{x-a} \frac{\omega(\varphi, h)}{(h+t)^{1+\alpha}} dt \leq C_1 \frac{\omega(\varphi, h)}{h^\alpha} + \int_h^\infty \frac{\omega(\varphi, h)}{(h+t)^{1+\alpha}} dt \leq C_2 \frac{\omega(\varphi, h)}{h^\alpha}$$

As for  $J_2$  we have  $|J_2| \leq \int_{x-a}^{x+h-a} (x+h-t)^{-1-\alpha} \omega(\varphi, x+h-t) dt \leq C \int_{x-a}^{x+h-a} t^{-1-\alpha} \omega(\varphi, t) dt$ . If  $x-a \leq h$  then

$$|J_2| \leq C \int_{x-a}^{2h} t^{-1-\alpha} \omega(\varphi, t) dt \leq C \int_0^{2h} t^{-1-\alpha} \omega(\varphi, t) dt \leq C_1 \int_0^h t^{-1-\alpha} \omega(\varphi, t) dt$$

If  $x-a \geq h$ , after the substitution  $t = \xi + x-a$  and taking into account the (almost) decreasing nature of the function  $t^{-1} \omega(\varphi, t)$ , we have

$$|J_2| \leq \int_0^h \frac{\omega(\varphi, x-a+\xi)}{(x-a+\xi)^{1+\alpha}} d\xi \leq C \frac{\omega(\varphi, h)}{h} \int_0^h \frac{d\xi}{(x-a+\xi)^\alpha} \leq C \frac{\omega(\varphi, h)}{h^\alpha}$$

Now let's estimate  $J_3$ . If  $x-a \leq h$  we have

$$|J_3| \leq \int_0^h \omega(\varphi, t) \frac{(t+h)^{1+\alpha} - t^{1+\alpha}}{t^{1+\alpha}(t+h)^{1+\alpha}} dt \leq C \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} \frac{h dt}{t+h} \leq C \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt$$

If  $x-a \geq h$ :

$$|J_3| \leq \int_0^h |\varphi(x) - \varphi(x-t)| \left| (h+t)^{-\alpha-1} - t^{-1-\alpha} \right| dt + \int_h^{x-a} |\varphi(x) - \varphi(x-t)| \left| (h+t)^{-\alpha-1} - t^{-1-\alpha} \right| dt \leq$$

$$\leq C_1 \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt + C_2 h \int_h^{x-a} \frac{\omega(\varphi, t)}{t^{2+\alpha}} dt \leq C_1 \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt + C_2 h \int_h^\infty \frac{\omega(\varphi, h)}{t^{2+\alpha}} dt \leq C \left[ \int_0^h \frac{\omega(\varphi, t)}{t^{1+\alpha}} dt + \frac{\omega(\varphi, h)}{h^\alpha} \right]$$

Gathering estimates for  $J_1, J_2, J_3$  we arrive at (11). The theorem is this proved.

### 3. Main result

We consider the mixed fractional Riemann-Liouville integral

$$(I_{a+,c+}^{\alpha,\beta} \varphi)(x, y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_c^y \frac{\varphi(t, s) dt ds}{(x-t)^{1-\alpha} (y-s)^{1-\beta}}, \quad x > a, y > c, 0 < \alpha, \beta < 1 \tag{13}$$

**Theorem 3.** Let  $\varphi(x, y) \in C(Q)$  and  $\varphi(x, y)|_{x=a, y=c} = 0$ . Then for the mixed fractional integral (13) we have estimates of the Zygmund type

$$\omega(I_{a+,c+}^{\alpha,\beta} \varphi; h, \eta) \leq C_1 h \eta \int_h^{b-a} \int_\eta^{d-c} \frac{\omega(\varphi; t, s)}{t^{2-\alpha} s^{2-\beta}} dt ds \tag{14}$$

$$\omega(I_{a+,c+}^{\alpha,\beta} \varphi; h, 0) \leq C_2 h \int_h^{b-a} t^{\alpha-2} \omega(\varphi; t, 0) dt, \quad \omega(I_{a+,c+}^{\alpha,\beta} \varphi; 0, \eta) \leq C_3 \eta \int_\eta^{d-c} s^{\beta-2} \omega(\varphi; 0, s) ds \tag{15}$$

We will not prove this theorem. Its proof can be found from [2], [3], [7], [13] and [14].

**Theorem 4.** Let  $\varphi(x, y)$  be continuous on  $Q$  and  $\varphi(x, y)|_{x=a, y=c} = 0$ . Then for mixed fractional derivative  $D_{a+,c+}^{\alpha,\beta} \varphi, 0 < \alpha, \beta < 1$  we have estimates of the Zygmund type

$$\omega(D_{a+,c+}^{\alpha,\beta} \varphi; h, \eta) \leq C_1 \int_0^h \int_0^\eta \frac{\omega(\varphi; t, s)}{t^{1+\alpha} s^{1+\beta}} dt ds \tag{16}$$

$$\omega(D_{a+,c+}^{\alpha,\beta} \varphi; h, 0) \leq C_2 \int_0^h t^{-\alpha-1} \omega(\varphi; t, 0) dt, \quad \omega(D_{a+,c+}^{\alpha,\beta} \varphi; 0, \eta) \leq C_3 \int_0^\eta s^{-\beta-1} \omega(\varphi; 0, s) ds \tag{17}$$

**Proof.** Using the identity (5), we represent the derivative (1)

$$(D_{a+,c+}^{\alpha,\beta} \varphi)(x, y) = \frac{1}{\Gamma(1-\alpha)\Gamma(1-\beta)} \left[ \frac{\varphi(a, c)}{(x-a)^\alpha (y-c)^\beta} + \frac{\psi_1(x)}{(y-c)^\beta} + \frac{\psi_2(y)}{(x-a)^\alpha} + \psi(x, y) \right]$$

where

$$\psi_1(x) = \frac{\varphi(x, c) - \varphi(a, c)}{(x-a)^\alpha} + \alpha \int_a^x \frac{\varphi(x, c) - \varphi(t, c)}{(x-t)^{\alpha+1}} dt, \quad \psi_2(y) = \frac{\varphi(a, y) - \varphi(a, c)}{(y-c)^\beta} + \beta \int_c^y \frac{\varphi(a, y) - \varphi(a, s)}{(y-s)^{\beta+1}} ds,$$

$$\psi(x, y) = \frac{\left( \Delta_{x-a, y-c}^{1,1} \varphi \right)(a, c)}{(x-a)^\alpha (y-c)^\beta} + \frac{\alpha}{(y-c)^\beta} \int_a^x \frac{\left( \Delta_{x-t, y-c}^{1,1} \varphi \right)(t, c)}{(x-t)^{1+\alpha}} dt + \frac{\beta}{(x-a)^\alpha} \int_c^y \frac{\left( \Delta_{x-a, y-s}^{1,1} \varphi \right)(a, s) ds}{(y-s)^{1+\beta}} +$$

$$+ \alpha\beta \int_a^x \int_c^y \frac{\left( \Delta_{x-t, y-s}^{1,1} \varphi \right) (t, s) dt ds}{(x-t)^{1+\alpha} (y-s)^{1+\beta}}.$$

According to the theorem  $\varphi(x, y)|_{x=a, y=c} = 0$ . Then  $\Psi_1(x) = 0$  and  $\Psi_2(y) = 0$ . We have

$$\left( D_{a+, c+}^{\alpha, \beta} \varphi \right) (x, y) = \frac{\Psi(x, y)}{\Gamma(1-\alpha)\Gamma(1-\beta)} = f(x, y) = f_1(x, y) + f_2(x, y) + f_3(x, y) + f_4(x, y).$$

We estimate each term separately.

Let  $h > 0, x, x+h \in [a, b]$ . We consider the differences:

$$\begin{aligned} |f_1(x+h, y) - f_1(x, y)| &\leq \frac{\left| \left( \Delta_{h, y-c}^{1,1} \varphi \right) (a, c) \right|}{(y-c)^\beta (x+h-a)^\alpha} + \frac{\left| \left( \Delta_{h, y-c}^{1,1} \varphi \right) (a, c) \right|}{(y-c)^\beta} \left| (x+h-a)^\alpha - (x-a)^\alpha \right| \leq \\ &\leq C_1 \left( \frac{\omega(\varphi; h, y-c)^{1,1}}{(y-c)^\beta (x+h-a)^\alpha} + \frac{\omega(\varphi; h, y-c)^{1,1}}{(y-c)^\beta} \left| (x+h-a)^\alpha - (x-a)^\alpha \right| \right), \\ |f_2(x+h, y) - f_2(x, y)| &\leq \int_a^x \frac{\left| \left( \Delta_{h, y-s}^{1,1} \varphi \right) (x, c) \right| dt}{(x+h-t)^{1+\alpha} (y-c)^\beta} + \int_{x-a}^{x+h-a} \frac{\left| \left( \Delta_{x+h-t, y-s}^{1,1} \varphi \right) (t, c) \right|}{(x+h-t)^{1+\alpha} (y-c)^\beta} dt + \\ &+ (y-c)^{-\beta} \int_a^x \left| \left( \Delta_{x-t, y-s}^{1,1} \varphi \right) (t, c) \right| \left| (x+h-t)^{-1-\alpha} - (x-t)^{-1-\alpha} \right| dt \leq \frac{C_2}{(y-c)^\beta} \left( \int_a^x \frac{\omega(\varphi; h, y-c)^{1,1}}{(x+h-t)^{1+\alpha}} dt + \right. \\ &\left. + \int_{x-a}^{x+h-a} \frac{\omega(\varphi; x+h-t, y-c)^{1,1}}{(x+h-t)^{1+\alpha}} dt + \int_a^x \omega(\varphi; x-t, y-c) \left| (x+h-t)^{-1-\alpha} - (x-t)^{-1-\alpha} \right| dt \right), \\ |f_3(x+h, y) - f_3(x, y)| &\leq \int_c^y \frac{\left| \left( \Delta_{h, y-s}^{1,1} \varphi \right) (a, s) \right| ds}{(x+h-a)^\alpha (y-s)^{1+\beta}} + \left| (x+h-a)^\alpha - (x-a)^\alpha \right| \int_c^y \frac{\left| \left( \Delta_{x-a, y-s}^{1,1} \varphi \right) (a, s) \right| ds}{(y-s)^{1+\beta}} \leq \\ &\leq C_3 \left( (x+h-a)^\alpha \int_c^y \frac{\omega(\varphi; h, y-s)^{1,1}}{(y-s)^{1+\beta}} ds + \left| (x+h-a)^\alpha - (x-a)^\alpha \right| \int_c^y \frac{\omega(\varphi; x-a, y-s)^{1,1}}{(y-s)^{1+\beta}} ds \right), \\ |f_4(x+h, y) - f_4(x, y)| &\leq \int_c^y \int_a^x \frac{\left| \left( \Delta_{h, y-s}^{1,1} \varphi \right) (x, s) \right| dt ds}{(x+h-t)^{1+\alpha} (y-s)^{1+\beta}} + \int_c^y \int_{x-a}^{x+h-a} \frac{\left| \left( \Delta_{x+h-t, y-s}^{1,1} \varphi \right) (t, s) \right|}{(x+h-t)^{1+\alpha} (y-s)^{1+\beta}} dt ds + \\ &+ \int_c^y \int_a^x \frac{\left| \left( \Delta_{x-t, y-s}^{1,1} \varphi \right) (t, s) \right|}{(y-s)^{1+\beta}} \left| (x+h-t)^{-1-\alpha} - (x-t)^{-1-\alpha} \right| dt ds \leq C_4 \left( \int_c^y \int_a^x \frac{\omega(\varphi; h, y-s)^{1,1}}{(x+h-t)^{1+\alpha} (y-c)^{1+\beta}} dt ds + \right. \\ &\left. + \int_c^y \int_{x-a}^{x+h-a} \frac{\omega(\varphi; x+h-t, y-s)^{1,1}}{(x+h-t)^{1+\alpha} (y-c)^{1+\beta}} dt ds + \int_c^y \int_a^x \frac{\omega(\varphi; x-t, y-s)^{1,1}}{(y-s)^{1+\beta}} \left| (x+h-t)^{-1-\alpha} - (x-t)^{-1-\alpha} \right| dt \right) \end{aligned}$$

Using estimations  $A_1, A_2, J_1, J_2, J_3$  of the proof of Theorem 2 and inequalities (2), (6), it's easily possible to receive an estimation

$$\omega(f; h, 0) \leq C_2 \int_0^h t^{-\alpha-1} \omega(\varphi; t, 0) dt$$

The estimate

$$\omega(f; 0, \eta) \leq C_3 \int_0^\eta s^{-\beta-1} \omega(\varphi; 0, s) ds$$

is symmetrical obtained.

Let  $h, \eta > 0$ ;  $x, x+h \in [a, b]$ ,  $y, y+\eta \in [c, d]$ . We consider the differences

$$\begin{aligned} \left( \Delta_{h,\eta}^{1,1} f_1 \right)(x, y) &= \frac{\left( \Delta_{h,\eta}^{1,1} \varphi \right)(x, y)}{(x+h-a)^\alpha (y+\eta-c)^\beta} + \frac{\left( \Delta_{h,y-c}^{1,1} \varphi \right)(x, c)}{(x+h-a)^\alpha} \left[ \frac{1}{(y-c)^\beta} - \frac{1}{(y+\eta-c)^\beta} \right] + \\ &+ \frac{\left( \Delta_{x-a,\eta}^{1,1} \varphi \right)(a, y)}{(y+\eta-c)^\beta} \left[ \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right] + \\ &+ \left( \Delta_{x-a,y-c}^{1,1} \varphi \right)(a, c) \left[ \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right] \left[ \frac{1}{(y-c)^\beta} - \frac{1}{(y+\eta-c)^\beta} \right] \\ \left( \Delta_{h,\eta}^{1,1} f_2 \right)(x, y) &= \frac{\beta}{(x+h-a)^\alpha} \int_{y-c}^{y+\eta-c} \frac{\left( \Delta_{h,y+\eta-s}^{1,1} \varphi \right)(x, s)}{(y+\eta-s)^{1+\beta}} ds + \frac{\beta}{(x+h-a)^\alpha} \int_0^y \frac{\left( \Delta_{h,\eta}^{1,1} \varphi \right)(x, y)}{(y+\eta-s)^{1+\beta}} ds + \\ &+ \beta \left[ \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right] \int_{y-c}^{y+\eta-c} \frac{\left( \Delta_{x-a,y+\eta-s}^{1,1} \varphi \right)(a, s)}{(y+\eta-s)^{1+\beta}} ds + \\ &+ \frac{\beta}{(x+h-a)^\alpha} \int_c^y \left( \Delta_{h,y-s}^{1,1} \varphi \right)(x, s) \left[ \frac{1}{(y-s)^{1+\beta}} - \frac{1}{(y+\eta-s)^{1+\beta}} \right] ds + \\ &+ \beta \left[ \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right] \int_c^y \frac{\left( \Delta_{x-a,\eta}^{1,1} \varphi \right)(a, y)}{(y+\eta-s)^{1+\beta}} ds + \\ &+ \beta \left[ \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right] \int_c^y \left( \Delta_{x-a,y-s}^{1,1} \varphi \right)(a, s) \left[ \frac{1}{(y-s)^{1+\beta}} - \frac{1}{(y+\eta-s)^{1+\beta}} \right] ds \\ \left( \Delta_{h,\eta}^{1,1} f_3 \right)(x, y) &= \frac{\alpha}{(y+\eta-c)^\beta} \int_{x-a}^{x+h-a} \frac{\left( \Delta_{x+h-t,\eta}^{1,1} \varphi \right)(t, y)}{(x+h-t)^{1+\alpha}} dt + \frac{\alpha}{(y+\eta-c)^\beta} \int_a^x \frac{\left( \Delta_{h,\eta}^{1,1} \varphi \right)(x, y)}{(x+h-t)^{1+\alpha}} dt + \\ &+ \alpha \left[ \frac{1}{(y-c)^\beta} - \frac{1}{(y+\eta-c)^\beta} \right] \int_{x-a}^{x+h-a} \frac{\left( \Delta_{x+h-t,y}^{1,1} \varphi \right)(t, c)}{(x+h-t)^{1+\alpha}} dt + \\ &+ \frac{\alpha}{(y+\eta-c)^\beta} \int_a^x \left( \Delta_{x-t,\eta}^{1,1} \varphi \right)(t, y) \left[ \frac{1}{(x-t)^{1+\alpha}} - \frac{1}{(x+h-t)^{1+\alpha}} \right] dt + \\ &+ \alpha \left[ \frac{1}{(y-c)^\beta} - \frac{1}{(y+\eta-c)^\beta} \right] \int_a^x \frac{\left( \Delta_{h,y-c}^{1,1} \varphi \right)(x, c)}{(x+h-t)^{1+\alpha}} dt + \\ &+ \alpha \left[ \frac{1}{(y-c)^\beta} - \frac{1}{(y+\eta-c)^\beta} \right] \int_a^x \left( \Delta_{x-t,y-c}^{1,1} \varphi \right)(t, c) \left[ \frac{1}{(x-t)^{1+\alpha}} - \frac{1}{(x+h-t)^{1+\alpha}} \right] dt \end{aligned}$$

$$\begin{aligned}
 \left(\Delta_{h,\eta}^{1,1} f_4\right)(x, y) &= \int_a^x \int_c^y \frac{\left(\Delta_{h,\eta}^{1,1} \varphi\right)(x, y) dt ds}{(x+h-t)^{1+\alpha}(y+\eta-s)^{1+\beta}} + \int_a^x \int_{y-c}^{y+\eta-c} \frac{\left(\Delta_{h,y+\eta-s}^{1,1} \varphi\right)(x, s) dt ds}{(x+h-t)^{1+\alpha}(y+\eta-s)^{1+\beta}} + \\
 &+ \int_a^x \int_c^y \frac{\left(\Delta_{h,y-s}^{1,1} \varphi\right)(x, s)}{(x+h-t)^{1+\alpha}} \left[ \frac{1}{(y-s)^{1+\beta}} - \frac{1}{(y+\eta-s)^{1+\beta}} \right] dt ds + \int_{x-a}^{x+h-a} \int_c^y \frac{\left(\Delta_{x+h-t,\eta}^{1,1} \varphi\right)(t, y) dt ds}{(x+h-t)^{1+\alpha}(y+\eta-s)^{1+\beta}} + \\
 &+ \int_{x-a}^{x+h-a} \int_{y-c}^{y+\eta-c} \frac{\left(\Delta_{x+h-t,y+\eta-s}^{1,1} \varphi\right)(t, s) dt ds}{(x+h-t)^{1+\alpha}(y+\eta-s)^{1+\beta}} + \int_{x-a}^{x+h-a} \int_c^y \frac{\left(\Delta_{x+h-t,y-s}^{1,1} \varphi\right)(t, s)}{(x+h-t)^{1+\alpha}} \left[ \frac{1}{(y-s)^{1+\beta}} - \frac{1}{(y+\eta-s)^{1+\beta}} \right] dt ds + \\
 &+ \int_a^x \int_c^y \frac{\left(\Delta_{x-t,\eta}^{1,1} \varphi\right)(t, y)}{(y+\eta-s)^{1+\beta}} \left[ \frac{1}{(x-t)^{1+\alpha}} - \frac{1}{(x+h-t)^{1+\alpha}} \right] dt ds + \\
 &+ \int_a^x \int_{y-c}^{y+\eta-c} \frac{\left(\Delta_{x-t,y+\eta-s}^{1,1} \varphi\right)(t, s)}{(y+\eta-s)^{1+\beta}} \left[ \frac{1}{(x-t)^{1+\alpha}} - \frac{1}{(x+h-t)^{1+\alpha}} \right] dt ds + \\
 &+ \int_a^x \int_c^y \frac{\left(\Delta_{x-t,y-s}^{1,1} \varphi\right)(t, s)}{(x-t)^{1+\alpha}} \left[ \frac{1}{(y-s)^{1+\beta}} - \frac{1}{(y+\eta-s)^{1+\beta}} \right] dt ds.
 \end{aligned}$$

The validity of these representations may be checked directly. We have

$$\begin{aligned}
 \left|\left(\Delta_{h,\eta}^{1,1} f_1\right)(x, y)\right| &\leq C_1 \left( \frac{\omega(\varphi; h, \eta)}{(x+h-a)^\alpha (y+\eta-c)^\beta} + \frac{\omega(\varphi; h, y-c)}{(x+h-a)^\alpha} \left| \frac{1}{(y-c)^\beta} - \frac{1}{(y+\eta-c)^\beta} \right| + \right. \\
 &+ \frac{\omega(\varphi; x-a, \eta)}{(y+\eta-c)^\beta} \left| \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right| + \\
 &\left. + \omega(\varphi; x-a, y-c) \left| \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right| \left| \frac{1}{(y-c)^\beta} - \frac{1}{(y+\eta-c)^\beta} \right| \right) \\
 \left|\left(\Delta_{h,\eta}^{1,1} f_2\right)(x, y)\right| &\leq C_2 \left( \frac{1}{(x+h-a)^\alpha} \int_{y-c}^{y+\eta-c} \frac{\omega(\varphi; h, y+\eta-s)}{(y+\eta-s)^{1+\beta}} ds + \frac{1}{(x+h-a)^\alpha} \int_0^y \frac{\omega(\varphi; h, \eta)}{(y+\eta-s)^{1+\beta}} ds + \right. \\
 &+ \left| \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right| \int_{y-c}^{y+\eta-c} \frac{\omega(\varphi; x-a, y+\eta-s)}{(y+\eta-s)^{1+\beta}} ds + \\
 &+ \frac{1}{(x+h-a)^\alpha} \int_c^{1,1} \omega(\varphi; h, y-s) \left| \frac{1}{(y-s)^{1+\beta}} - \frac{1}{(y+\eta-s)^{1+\beta}} \right| ds + \\
 &+ \left| \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right| \int_c^y \frac{\omega(\varphi; x-a, \eta)}{(y+\eta-s)^{1+\beta}} ds + \\
 &+ \left| \frac{1}{(x-a)^\alpha} - \frac{1}{(x+h-a)^\alpha} \right| \int_c^{1,1} \omega(\varphi; x-a, y-s) \left| \frac{1}{(y-s)^{1+\beta}} - \frac{1}{(y+\eta-s)^{1+\beta}} \right| ds \Big) \\
 \left|\left(\Delta_{h,\eta}^{1,1} f_3\right)(x, y)\right| &\leq C_3 \left( \frac{1}{(y+\eta-c)^\beta} \int_{x-a}^{x+h-a} \frac{\omega(\varphi; x+h-t, \eta)}{(x+h-t)^{1+\alpha}} dt + \frac{1}{(y+\eta-c)^\beta} \int_a^x \frac{\omega(\varphi; h, \eta)}{(x+h-t)^{1+\alpha}} dt + \right. \\
 &+ \left. \left| \frac{1}{(y-c)^\beta} - \frac{1}{(y+\eta-c)^\beta} \right| \int_{x-a}^{x+h-a} \frac{\omega(\varphi; x+h-t, y-s)}{(x+h-t)^{1+\alpha}} dt + \right.
 \end{aligned}$$



$$\begin{aligned}
 & + \frac{1}{(y + \eta - c)^\beta} \int_a^{x,1,1} \omega(\varphi; x - t, \eta) \left| \frac{1}{(x - t)^{1+\alpha}} - \frac{1}{(x + h - t)^{1+\alpha}} \right| dt + \\
 & + \left| \frac{1}{(y - c)^\beta} - \frac{1}{(y + \eta - c)^\beta} \right| \int_a^{x,1,1} \frac{\omega(\varphi; h, y - c)}{(x + h - t)^{1+\alpha}} dt + \\
 & + \left( \frac{1}{(y - c)^\beta} - \frac{1}{(y + \eta - c)^\beta} \int_a^{x,1,1} \omega(\varphi; x - t, y - c) \left| \frac{1}{(x - t)^{1+\alpha}} - \frac{1}{(x + h - t)^{1+\alpha}} \right| dt \right) \\
 & \left| \left( \Delta_{h,\eta} f_4 \right)(x, y) \right| \leq C_4 \left( \int_a^x \int_c^y \frac{\omega(\varphi; h, \eta) dt ds}{(x + h - t)^{1+\alpha} (y + \eta - s)^{1+\beta}} + \int_a^x \int_{y-c}^{y+\eta-c} \frac{\omega(\varphi; h, y + \eta - s) dt ds}{(x + h - t)^{1+\alpha} (y + \eta - s)^{1+\beta}} + \right. \\
 & + \int_a^x \int_c^y \frac{\omega(\varphi; h, y - s)}{(x + h - t)^{1+\alpha}} \left| \frac{1}{(y - s)^{1+\beta}} - \frac{1}{(y + \eta - s)^{1+\beta}} \right| dt ds + \int_{x-a}^{x+h-a} \int_c^y \frac{\omega(\varphi; x + h - t, \eta) dt ds}{(x + h - t)^{1+\alpha} (y + \eta - s)^{1+\beta}} + \\
 & + \int_{x-a}^{x+h-a} \int_{y-c}^{y+\eta-c} \frac{\omega(\varphi; x + h - t, y + \eta - s) dt ds}{(x + h - t)^{1+\alpha} (y + \eta - s)^{1+\beta}} + \\
 & + \int_{x-a}^{x+h-a} \int_c^{y,1,1} \frac{\omega(\varphi; x + h - t, y - s)}{(x + h - t)^{1+\alpha}} \left| \frac{1}{(y - s)^{1+\beta}} - \frac{1}{(y + \eta - s)^{1+\beta}} \right| dt ds + \\
 & + \int_a^x \int_c^y \frac{\omega(\varphi; x - t, \eta)}{(y + \eta - s)^{1+\beta}} \left| \frac{1}{(x - t)^{1+\alpha}} - \frac{1}{(x + h - t)^{1+\alpha}} \right| dt ds + \\
 & + \int_a^x \int_{y-c}^{y+\eta-c} \frac{\omega(\varphi; x - t, y + \eta - s)}{(y + \eta - s)^{1+\beta}} \left| \frac{1}{(x - t)^{1+\alpha}} - \frac{1}{(x + h - t)^{1+\alpha}} \right| dt ds + \\
 & + \int_a^x \int_c^{y,1,1} \omega(\varphi; x - t, y - s) \left| \frac{1}{(x - t)^{1+\alpha}} - \frac{1}{(x + h - t)^{1+\alpha}} \right| \left| \frac{1}{(y - s)^{1+\beta}} - \frac{1}{(y + \eta - s)^{1+\beta}} \right| dt ds \Big).
 \end{aligned}$$

After which every term is estimated in the standard way and we get

$$\omega(f; h, \eta) \leq C_1 \int_0^h \int_0^\eta \frac{\omega(\varphi; t, s)}{t^{1+\alpha} s^{1+\beta}} dt ds$$

This completes the proof.

**4. References**

- Samko SG, Kilbas AA, Marichev OI. Fractional Integrals and Derivatives. Theory and Applications. Gordon and Breach. Sci. Publ., N. York - London, 1993, 1012 pp. (Russian Ed. - Fractional Interals and Derivatives and Some of Their Applications, Nauka i Tekhnika, Minsk, 1987.
- Mamatov T. Weighted Zygmund estimates for mixed fractional integration. Case Studies Journal. 2018; 7(5):82-88
- Mamatov T. Mixed Fractional Integration In Mixed Weighted Generalized Hölder Spaces. Case Studies Journal. 2018; 7(6):61-68
- Mamatov T. Mixed Fractional Integration Operators in Mixed Weighted Hölder Spaces. Monograph. LAPLAMBERT Academic Publishing. 73.
- Mamatov T. Mixed Fractional Integro-Differentiation Operators in Hölder Spaces. The latest research in modren science: experience, traditions and innovations. Proceedings of the VII International Scientific Conference. Section I. North Charleston, SC, USA, 2018, 6-9
- Mamatov T, Rayimov D, Elmurodov M. Mixed Fractioanl Differentiation Operators in Hölder Spaces. Journal of Multidisciplinary Engineering Science and Technology (JMEST), 2019; 6(4):9855-9857.
- Mamatov T. Fractional integration operators in mixed weighted generalized Hölder spaces of function of two variables defined by mixed modulus of continuity. "Journal of Mathematical Methods in Engineering" Auctores Publishing. 2019; 1(1):1-16. [www.auctoresonline.org](http://www.auctoresonline.org). (DOI:10.31579/jmme.2019/004) 2019
- Mamatov T. "Mapping Properties of Mixed Fractional Integro-Differentiation in Hölder Spaces", Journal of Concrete and Applicable Mathematics (JCAAM). 2014; 12(3-4):272-290
- Mamatov T. Mapping Properties of Mixed Fractional Differentiation Operators in Hölder Spaces Defined by Usual Hölder Condition, Journal of Computer Science & Computational Mathematics, 2019; 9(2). DOI: 10.20967/jcscm.2019.02.003

10. Mamatov T, Homidov F, Rayimov D. On Isomorphism Implemented by Mixed Fractional Integrals In Hölder Spaces, *International Journal of Development Research*. 2019; 9(5):27720-27730.
11. Mamatov T. Composition of mixed Riemann-Liouville fractional integral and mixed fractional derivative. *Journal of Global Research in Mathematical Archives*. 2019; 6(11):23-32. [Online]. Available: <http://www.jgrma.info>.
12. Mamatov T, Rahimov D, Some properties of mixed fractional integro-differentiation operators in Hölder spaces. *Journal of Global Research in Mathematical Archives*. 2019; 6(11):13-22. [Online]. Available: <http://www.jgrma.info>.
13. Mamatov T, Homidov F. Zigmund type estimates for mixed fractional integrals of the Volterra convolution type. *Chronos Journal*. 11(37):82-86. [www.chronos-journal.ru](http://www.chronos-journal.ru)
14. Mamatov T, Mustafoev N. Non-Weighted Zygmund Type Estimates for The Volterra Convolution Type. *Impact Factor 3.582 Case Studies Journal*. 2019; 8(11):119-122 ISSN (2305-509X). <http://www.casestudiesjournal.com>
15. Mamatov T, Umarov A, Rustamova L. Mixed Fractional Differentiation Operators in Mixed Weighted Hölder Spaces. *Impact Factor 3.582 Case Studies Journal*. 2019; 8(11):113-118 ISSN (2305-509X) <http://www.casestudiesjournal.com>.