



Journal of Mathematical Problems, Equations and Statistics

E-ISSN: 2709-9407

P-ISSN: 2709-9393

JMPES 2021; 2(2): 43-47

© 2021 JMPES

www.mathematicaljournal.com

Received: 17-04-2021

Accepted: 20-06-2021

Tulkin Mamatov

Department of Higher
mathematics, Bukhara
Technological Institute of
Engineering, Bukhara,
Uzbekistan

Hardy-little wood-type theorem for mixed fractional integrals in weighted holder spaces

Tulkin Mamatov

Abstract

We study mixed Riemann-Liouville fractional integration operators and mixed fractional derivative in Marchaud form of function of two variables in Hölder spaces of different orders in each variables. The obtained are results generalized to the case of Hölder spaces with power weight.

Keywords: Functions of two variables, fractional derivative of Marchaud form, mixed fractional derivative, weight, mixed fractional integral, Hölder space.

Introduction

In 1928 H.G. Hardy and J.E. little wood ^[1] (see ^[15], Theorem 3.1 and 3.2) showed that the fractional integral

$$(I_{0+}^{\alpha} \varphi)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x \frac{\varphi(t) dt}{(x-t)^{1-\alpha}}, \quad 0 < x < 1$$

of order $\alpha \in (0,1)$ improves the Hölder behavior of its density exactly by the order α . More exactly, these operators establish an isomorphism between the spaces $H_0^{\lambda}([0,1])$ and $H_0^{\lambda+\alpha}([0,1])$ under the condition $\alpha + \lambda$. This result was extended in many directions: to the case of Hölder spaces with power weight ^[14] to the case of generalized Hölder spaces with characteristics from the Bari-Stechkin class ^[12, 13]; to the case of more general weights ^[15, 16], etc. Different proofs were suggested in ^[2, 3], where the case of complex fractional orders was also considered the shortest proof being given in ^[2].

In the multidimensional case, the statement about the properties of a map in Hölder spaces for a mixed fractional Riemann - Liouville integral was studied in ^[4, 11].

As is known, the Riemann-Liouville fractional integration operator establishes an isomorphism between weighted Hölder spaces for functions one variable. But for functions two variable were not studied. This paper is aimed to fill in this gap. We study mixed Riemann-Liouville fractional integration operators and mixed fractional derivative in Marchaud form of function of two variables in Hölder spaces of different orders in each variables. The obtained results generalized to the case of Hölder spaces with power weight of functions of two variables.

Mixed Riemann-Liouville fractional integrals of order (α, β)

$$(I_{0+,0+}^{\alpha,\beta} \varphi)(x, y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^x \int_0^y \frac{\varphi(t, s) dt ds}{(x-t)^{1-\alpha} (y-s)^{1-\beta}}, \quad (1)$$

$x > 0, y > 0$.

Mixed fractional derivatives form Marchaud of order (α, β)

Correspondence

Tulkin Mamatov

Department of Higher
mathematics, Bukhara
Technological Institute of
Engineering, Bukhara,
Uzbekistan

$$\begin{aligned} (D_{0+,0+}^{\alpha,\beta} \varphi)(x,y) &= \frac{\varphi(x,y)}{\Gamma(1-\alpha)\Gamma(1-\beta)} x^{-\alpha} y^{-\beta} + \\ &+ \frac{\alpha\beta}{\Gamma(\alpha)\Gamma(\beta)} \int_0^x \int_0^y \frac{\varphi(x,y) - \varphi(t,s)}{(x-t)^{1+\alpha} (y-s)^{1+\beta}} dt ds, \quad x > 0, y > 0 \end{aligned} \tag{2}$$

Consider the operators in a rectangle

$$Q = \{(x, y) : 0 < x < b, 0 < y < d\}$$

For a continuous function $\varphi(x, y)$ on R^2 we introduce the notation

$$\begin{aligned} \left(\Delta_h^{1,0} \varphi\right)(x,y) &= \varphi(x+h,y) - \varphi(x,y), \\ \left(\Delta_\eta^{0,1} \varphi\right)(x,y) &= \varphi(x,y+\eta) - \varphi(x,y), \\ \left(\Delta_{h,\eta}^{1,1} \varphi\right)(x,y) &= \varphi(x+h,y+\eta) - \varphi(x+h,y) - \varphi(x,y+\eta) + \varphi(x,y). \end{aligned}$$

So that

$$\begin{aligned} \varphi(x+h,y+\eta) &= \left(\Delta_{h,\eta}^{1,1} \varphi\right)(x,y) + \left(\Delta_\eta^{0,1} \varphi\right)(x,y) + \\ &+ \left(\Delta_h^{1,0} \varphi\right)(x,y) + \varphi(x,y). \end{aligned} \tag{3}$$

Everywhere in the sequel by C, C_1, C_2 etc we denote positive constants which may have different values in different occurrences and even in the same line.

Definition 1. Let $\lambda, \gamma \in (0,1]$. We say that $\varphi(x, y) \in H^{\lambda,\gamma}(Q)$, if

$$|\varphi(x_1, y_1) - \varphi(x_2, y_2)| \leq C_1 |x_1 - x_2|^\lambda + C_2 |y_1 - y_2|^\gamma \tag{4}$$

For all $(x_i, y_i) \in Q, i = 1, 2$.

Condition (4) is equivalent to the couple of the separate conditions

$$\left| \left(\Delta_h^{1,0} \varphi\right)(x,y) \right| \leq C_1 |h|^\lambda, \quad \left| \left(\Delta_\eta^{0,1} \varphi\right)(x,y) \right| \leq C_2 |\eta|^\gamma$$

Uniform with respect to another variable.

By $H_0^{\lambda,\gamma}(Q)$ we define a subspace of functions $\varphi \in H^{\lambda,\gamma}(Q)$, vanishing at the boundaries $x = 0$ and $y = 0$ of Q .

Let $\lambda = 0$ and/or $\gamma = 0$. We put $H^{0,0}(Q) = L^\infty(Q)$ and

$$H^{\lambda,0}(Q) = \left\{ \varphi \in L^\infty(Q) : \left| \left(\Delta_h^{1,0} \varphi\right)(x,y) \right| \leq C_1 |h|^\lambda \right\}, \lambda \in (0,1]$$

$$H^{0,\gamma}(Q) = \left\{ \varphi \in L^\infty(Q) : \left| \left(\Delta_\eta^{0,1} \varphi\right)(x,y) \right| \leq C_2 |\eta|^\gamma \right\}, \gamma \in (0,1].$$

Definition 2. We say that $\varphi(x, y) \in \tilde{H}^{\lambda, \gamma}(Q)$, where $\lambda, \gamma \in (0, 1]$, if

$$\varphi \in H^{\lambda, \gamma}(Q) \text{ And } \left| \left(\Delta_{h, \eta}^{1,1} \varphi \right) (x, y) \right| \leq C_3 |h|^\lambda |\eta|^\gamma.$$

We say that $\varphi \in \tilde{H}_0^{\lambda, \gamma}(Q)$, if $\varphi \in H^{\lambda, \gamma}(Q)$ any $\varphi(0, y) \equiv \varphi(x, 0) \equiv 0$.

These spaces become Banach spaces under the standard definition of the norms:

$$\begin{aligned} \|\varphi\|_{H^{\lambda, \gamma}} &:= \|\varphi\|_{C(Q)} + \sup_{\substack{x, x+h \in [0, b] \\ y \in [0, d]}} \frac{\left| \left(\Delta_h^{1,0} \varphi \right) (x, y) \right|}{|h|^\lambda} + \sup_{\substack{y, y+\eta \in [0, b] \\ x \in [0, b]}} \frac{\left| \left(\Delta_\eta^{0,1} \varphi \right) (x, y) \right|}{|\eta|^\gamma}, \\ \|\varphi\|_{\tilde{H}^{\lambda, \gamma}} &:= \|\varphi\|_{H^{\lambda, \gamma}} + \sup_{\substack{x, x+h \in [0, b] \\ y, y+\eta \in [0, d]}} \frac{\left| \left(\Delta_{h, \eta}^{1,1} \varphi \right) (x, y) \right|}{|h|^\lambda |\eta|^\gamma}. \end{aligned}$$

Note that

$$\varphi(x, y) \in H^{\lambda, \gamma}(Q) \Rightarrow \left| \left(\Delta_{h, \eta}^{1,1} \varphi \right) (x, y) \right| \leq C_\theta |h|^{\theta\lambda} |\eta|^{(1-\theta)\gamma} \tag{5}$$

For any $\theta \in [0, 1]$, where $C_\theta = 2C_1^\theta C_2^{1-\theta}$.

So that

$$\bigcap_{0 \leq \theta \leq 1} \tilde{H}^{\lambda\theta, (1-\theta)\gamma}(Q) \lrcorner H^{\lambda, \gamma}(Q) \lrcorner \tilde{H}^{\lambda, \gamma}(Q) \tag{6}$$

Where \lrcorner stands for the continuous embedding.

The norm for $\bigcap_{0 \leq \theta \leq 1} \tilde{H}^{\lambda\theta, (1-\theta)\gamma}(Q)$ is introduced as the maximum in θ of norms for $\tilde{H}^{\lambda\theta, (1-\theta)\gamma}(Q)$. Since $\theta \in [0, 1]$ is

arbitrary, it is not hard to see that the inequality in (5) is equivalent (up to the constant factor C) to

$$\left| \left(\Delta_{h, \eta}^{1,1} \varphi \right) (x, y) \right| \leq C \min \{ |h|^\lambda, |\eta|^\gamma \}. \tag{7}$$

We will also make use of the following weighted spaces. Let $\rho(x, y)$ be a non-negative function on Q (we will only deal with degenerate weights

$$\rho(x, y) = \rho_1(x)\rho_2(y).$$

Definition 3. By $H^{\lambda, \gamma}(Q, \rho)$ and $\tilde{H}^{\lambda, \gamma}(Q, \rho)$ we denote the spaces of functions $\varphi(x, y)$ such that $\rho\varphi \in H^{\lambda, \gamma}(Q)$ and $\rho\varphi \in \tilde{H}^{\lambda, \gamma}(Q)$ respectively, equipped with the norms

$$\|\varphi\|_{H^{\lambda, \gamma}(Q, \rho)} = \|\rho\varphi\|_{H^{\lambda, \gamma}(Q)} \text{ and } \|\varphi\|_{\tilde{H}^{\lambda, \gamma}(Q, \rho)} = \|\rho\varphi\|_{\tilde{H}^{\lambda, \gamma}(Q)}.$$

By $H_0^{\lambda,\gamma}(Q, \rho)$ and $\tilde{H}_0^{\lambda,\gamma}(Q, \rho)$ we denote the corresponding subspaces of functions φ such that

$$\rho\varphi|_{x=0} = \rho\varphi|_{y=0} \equiv 0.$$

Below we follow some technical estimations suggested in [2] for the case of one-dimensional Riemann-Liouville fractional integrals. We denote

$$B(x, y; t, s) = \frac{\rho(x, y) - \rho(t, s)}{\rho(t, s)(x-t)^\delta (y-s)^\sigma} \tag{8}$$

Where

$$0 < \delta, \sigma < 2, a < t < x < b, 0 < s < y < d.$$

Main Result

The definition in the Marchaud form may be used for all $-1 < \alpha, \beta < 1$: if $\alpha, \beta > 0$ (2) gives the mixed fractional derivative, if $\alpha, \beta < 0$, it is mixed fractional integral.

Let $\rho(x, y) = \rho(x)\rho(y)$ be the weight function and put

$$\psi(x, y) = \rho(x, y)\varphi(x, y) \quad \varphi \in \tilde{H}_0^{\lambda,\gamma}(Q; \rho)$$

Evidently $\psi \in \tilde{H}^{\lambda,\gamma}(Q)$ and $\psi(x, y)|_{x=0, y=0} = 0$ It is easy to see that

$$(\rho J_{0+,0+}^{\alpha,\beta} \varphi)(x, y) = (J_{0+,0+}^{\alpha,\beta} \psi)(x, y) + C(K_{0+,0+}^{\alpha,\beta} \psi)(x, y), \tag{9}$$

Where $-1 < \alpha, \beta < 1$, C =constant and

$$(K_{0+,0+}^{\alpha,\beta} \psi)(x, y) = \int_0^x \int_0^y B(x, y; t, s) \psi(t, s) dt ds,$$

So that in (9) we have the mixed fractional integral if $0 \leq \alpha, \beta < 1$ and the mixed fractional derivative if

$$-1 < \alpha, \beta \leq 0.$$

The representation (9) for the fractional (integral) derivative shows that the estimate for the continuity modulus in the weighted case reduces to two simpler estimates:

- 1) The known non-weighted estimate of Hardy-Littlewood for mixed fractional integral (see [4]) and mixed fractional derivative (see [7]); in the case weighted estimated of Hardy-Littlewood for mixed fractional integral (see [4]);
- 2) The estimate of the second term in (9), which is the mixed fractional derivative. It is main part of the job.

Theorem 2. Let $0 < \alpha, \beta < 1, 0 < \lambda, \gamma < 1, \rho(x, y) = x^\mu y^\nu$

And let $\alpha + \lambda < 1, \beta + \gamma < 1$ and $|\psi(x, y)| \leq Cx^{\lambda+\alpha} y^{\gamma+\beta}$

With $\mu < 1 + \lambda, \nu < 1 + \gamma$. Then the operator $K_{0+,0+}^{\alpha,\beta}$ is bounded from the space

$$\tilde{H}_0^{\lambda,\gamma}(\rho) \text{ into } \tilde{H}_0^{\lambda+\alpha,\gamma+\beta}(\rho).$$

Proof. We don't proof this theorem. The proof of this theorem can be seen in [4].

Theorem 3: Let $0 < \alpha, \beta < 1, 0 < \lambda, \gamma < 1, \rho(x, y) = x^\mu y^\nu$ and let $\alpha + \lambda < 1, \beta + \gamma < 1$

and $|\psi(x, y)| \leq Cx^\lambda y^\gamma$

With $\mu < 1 + \lambda, \nu < 1 + \gamma$. Then the operator $K_{0+,0+}^{-\alpha,-\beta}$ is bounded from the space $\tilde{H}_0^{\lambda,\gamma}(\rho)$ into $\tilde{H}_0^{\lambda-\alpha,\gamma-\beta}(\rho)$.

Proof. The proof of this theorem can be seen in [8].

Theorem 4. Let $\rho(x, y)$ have the form $\rho(x, y) = x^\mu y^\nu$ with $\mu < 1 + \lambda, \nu < 1 + \gamma$ and let $0 < \alpha, \beta < 1, 0 < \lambda, \gamma < 1, \lambda + \alpha < 1, \gamma + \beta < 1$. Then the operator $J_{0+,0+}^{\alpha,\beta}$ establishes an isomorphism between the spaces $\tilde{H}_0^{\lambda,\gamma}(\rho)$ and $\tilde{H}_0^{\lambda+\alpha,\beta+\gamma}(\rho)$.

Proof. We should consider, as usual the following three parts of the proof:

1. Action of the mixed fractional integral operator from the space $\tilde{H}_0^{\lambda,\gamma}(\rho)$ to the space $\tilde{H}_0^{\lambda+\alpha,\beta+\gamma}(\rho)$;
2. Action of the mixed fractional differentiation operator from the space $\tilde{H}_0^{\lambda+\alpha,\beta+\gamma}(\rho)$ to the space $\tilde{H}_0^{\lambda,\gamma}(\rho)$;
3. The possibility to represent any function $f(x, y) \in \tilde{H}_0^{\lambda+\alpha,\beta+\gamma}(\rho)$ as $(J_{0+,0+}^{\alpha,\beta}\varphi)(x, y)$ with the density in $\tilde{H}_0^{\lambda,\gamma}(\rho)$.

Because of $|\psi(t, s)| \leq Ct^{\lambda+\alpha} s^{\gamma+\beta}, |\psi(t, s) - \psi(x, 0)| \leq C(t-x)^{\lambda+\alpha} s^{\gamma+\beta}$ the parts 1) -2) are covered by Theorems 2 and 3. The part 3) is treated in the standard way in case $0 < \alpha, \beta < 1$ by using the possibility of similar representation with the density from $L_{\bar{p}}(\mathbb{R}^2), \bar{p} = (p_1, p_2)$. See [15] Theorem 24.4.

References

1. Hardy HG, Littlewood JE. Some properties of fractional integrals. I. Math. Z 1928;27(4):565-606.
2. Karapetiants NK, Shankishvili LD. A shot proof of Hardy- Littlewood – type theorem for fractional integrals in weighted Hölder spaces. Fract. Calc. Appl. Anal 1999;2(2):117-192.
3. Karapetiants NK, Shankishvili LD. Fractional integro-differentiation of the complex order in generalized Hölder spaces $H_0^\omega([0,1], \rho)$. Integral Transforms Spec. Funct 2002;13(3):199-209.
4. Mamatov T, Samko S. Mixed fractional integration operators in mixed weighted Hölder spaces. Fract. Calc. Appl. Anal 2010;13(3):245-259.
5. Mamatov T. Mixed fractional integration operators in mixed weighted generalized Hölder spaces. Case Studies Journal 2018;7(6):61-68.
6. Mamatov T, Mapping properties of mixed fractional differentiation operators in Hölder spaces defined by usual Hölder condition. Journal of Computer Science and Computational Mathematics 2019;2(9):29-34.
7. T. Mamatov, Rayimov D., Some properties of mixed fractional integro-differentiation operators in Hölder spaces. Journal of Global Research in Mathematical Archives 2019;6(11):13-22.
8. Mamatov T, Rayimov D, Homidov F. On isomorphism implemented by mixed fractional integrals in Hölder spaces. International Journal of Development Research (IJDR) ISSN: 2230-9926., 2019;9:27720-27730.
9. Mamatov T. Fractional integration operators in mixed weighted generalized Hölder spaces of function of two variables defined by mixed modulus of continuity. “Journal of Mathematical Methods in Engineering” Auctores Publishing, vol. 2019, 1(1)-004, 1-16.
10. Mamatov T. Mixed fractional integration operators in mixed weighted Hölder spaces// (Monograph Online Sources), LAP LAMBERT Academic Publishing, Beau Bassin 2018, 73.
11. Mamatov T, Operator of Volterra convolution type in generalized Hölder spaces. Poincare Journal of Analysis & Applications 2020;7(2):275-288.
12. Kh M Murdaev, Samko SG. Weighted estimates of continuity modulus of fractional integrals of function having a prescribed continuity modulus with weight. Deponiert in VINITI, Moscow 1986;3351-B:42.
13. Kh. M. Murdaev, Samko SG, Mapping properties of fractional integro-differentiation in weighted generalized Hölder spaces $H_0^\omega(\rho)$ with the weight $\rho(x) = (x-a)^\mu (b-x)^\nu$ and given continuity modulus. Deponiert in VINITI, Moscow 1986;3350-B:25.
14. Rubin BS, Fractional integrals in Hölder spaces with weight and operators of potential type. Izv. Akad. Nauk Armyan. SSR Ser. Math 1974;9(4):308-324.
15. Samko SG, Kilbas AA, Marichev OI. Fractional Integrals and Derivatives (Book style). Theory and Applications, Gordon and Breach. Sci. Publ. N. York-London 1993, 1012.
16. Samko SG, Mussalavaeva Z. Fractional type operators in weighted generalized Hölder spaces. Proc. Georgian Acad. Sci. Mathem 1993;1(5):601-626.