



# Journal of Mathematical Problems, Equations and Statistics

E-ISSN: 2709-9407

P-ISSN: 2709-9393

JMPES 2020; 1(2): 63-68

© 2020 JMPES

[www.mathematicaljournal.com](http://www.mathematicaljournal.com)

Received: 22-04-2020

Accepted: 25-05-2020

**Anshu Sharma**Department of Mathematics,  
Govt. Mam PG College, Jammu,  
Jammu & Kashmir, India

## Compact weighted type composition operators from $Q_K(p, q)$ spaces to Bloch-type spaces

**Anshu Sharma****Abstract**

For a non-negative integer  $n$ ; the generalized weighted composition operator  $D_{\varphi, \psi}^n$  is defined by  $D_{\varphi, \psi}^n f = \psi \cdot (f^{(n)} \circ \varphi)$ ,  $f \in H(\mathbb{D})$ , where  $\psi$  be holomorphic map of the open unit disk  $\mathbb{D}$ ,  $\varphi$  a holomorphic self-map  $\mathbb{D}$  and  $H(\mathbb{D})$  be the space of holomorphic functions on  $\mathbb{D}$ . In this paper, we characterize compactness of  $D_{\varphi, \psi}^n$  from  $Q_K(p, q)$  spaces to Bloch-type spaces.

**Keywords:** Compact weighted, Bloch-type spaces,  $Q_K(p, q)$ **Introduction**

Let  $\mathbb{D}$  be the open unit disk in the complex plane  $\mathbb{C}$ ,  $\partial\mathbb{D}$  its boundary,  $dA(z)$  the normalized area measure on  $\mathbb{D}$  (i.e.  $A(\mathbb{D}) = 1$ ),  $H(\mathbb{D})$  the class of all holomorphic functions on  $\mathbb{D}$ . Let  $0 < p < \infty$ ,  $-2 < q < \infty$  and  $K: [0, \infty) \rightarrow [0, \infty)$  a non-decreasing continuous function. A function  $f \in H(\mathbb{D})$  is in  $Q_K(p, q)$  if

$$M(f) = \left\{ \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^q K(g(z, a)) dA(z) \right\}^{1/p} < \infty,$$

Where

$$g(z, a) = \log \left| \frac{1 - \bar{a}z}{a - z} \right| = \log \frac{1}{|\eta_a(z)|}$$

is the Green function in  $\mathbb{D}$  with logarithmic singularity at  $a$ .

2000 Mathematics Subject Classification. Primary 47B33; Secondary 30H05, 46E15.

Key words and phrases. Compact weighted composition operator,  $Q_K(p, q)$  space,  $F(p, q, s)$  space, Bloch-type space.

Throughout this paper, we assume that

$$\int_0^1 (1 - r^2)^q K(-\log r) r dr < \infty, \quad (1)$$

since otherwise  $Q_K(p, q)$  consists only of constant functions (see [19]). For  $1 \leq p < \infty$ ,  $Q_K(p, q)$  is a Banach space with respect to the norm  $\|f\|_{Q_K(p, q)} = |f(0)| + M(f)$ . If  $K(x) = x^s$ ,  $s \geq 0$ , then the spaces  $Q_K(p, q)$  reduces to  $F(p, q, s)$  which was introduced by R. Zhao in [18], is known as general family of function spaces. Let  $\omega$  be a strictly positive continuous function on  $\mathbb{D}$ . If  $\omega(z) = \omega(|z|)$  for every  $z \in \mathbb{D}$ , we call it a radial weight. A radial weight  $\omega$  is called typical if it is non-increasing with respect to  $|z|$  and  $\omega(z) \rightarrow 0$  as  $|z| \rightarrow 1$ . For a typical weight  $\omega$ , the Bloch-type space  $\mathcal{B}_\omega$  on  $\mathbb{D}$  is a Banach space of  $f \in H(\mathbb{D})$  such that

$$\|f\|_{\mathcal{B}_\omega} = |f(0)| + \sup_{z \in \mathbb{D}} \omega(z) |f'(z)|.$$

The weighted spaces of controlled growth  $\mathcal{A}_\omega$  consists of  $f \in H(\mathbb{D})$  such that  $\|f\|_{\mathcal{A}_\omega} = \sup_{z \in \mathbb{D}} \omega(z) |f(z)| < \infty$ .

Let  $\psi \in H(\mathbb{D})$  and  $\varphi$  be a holomorphic map of  $\mathbb{D}$ . For a non-negative integer  $n$ , we define a linear operator  $D_{\varphi, \psi}^n$  as follows:

**Corresponding Author:****Anshu Sharma**Department of Mathematics,  
Govt. Mam PG College, Jammu,  
Jammu & Kashmir, India

$$D_{\varphi,\psi}^n f = \psi \cdot (f^{(n)} \circ \varphi), \quad f \in H(\mathbb{D}).$$

For example, if  $n = 0$  and  $\psi \equiv 1$ , then we obtain the composition operator  $C_\varphi$  induced by  $\varphi$ , defined as  $C_\varphi f = f \circ \varphi$ ,  $f \in H(\mathbb{D})$ . If  $\psi = 1$  and  $\varphi(z) = z$ , then  $D_{p,\psi}^n = D^n$ , the differentiation operator defined as  $D^n f = f^{(n)}$ . If  $n = 0$ , then we get the weighted composition operator  $\psi C_\varphi$  defined as  $\psi C_\varphi f = \psi \cdot (f \circ \varphi)$ .

For more about operators of the type  $D_{p,\psi}^n$  we refer [1-17]. In this paper, we characterize boundedness and compactness of  $D_{\varphi,\psi}^n$  from  $Q_K(p, q)$  spaces to Bloch-type spaces.

Throughout this paper constants are denoted by  $C$ , they are positive and not necessarily the same at each occurrence. The notation  $A \asymp B$  means that  $B \lesssim A \lesssim B$ , where  $A \lesssim B$  means that there is a positive constant  $C$  such that  $A \leq CB$ .

### Compactness of $D_{\varphi,\psi}^n$

In this section, we characterize compactness of  $D_{\varphi,\psi}^n$  from  $Q_K(p, q)$  spaces to Bloch-type spaces.

For the purpose we need several lemmas. The first among these can be found in [19].

**Lemma 1.** Let  $0 < p < \infty$ ,  $-2 < q < \infty$ ,  $\alpha = (2 + q)/p$ ,  $K: [0, \infty) \rightarrow [0, \infty)$  a non-decreasing continuous function such that (1) holds and  $\in Q_K(p, q)$ . Then (a)  $Q_K(p, q) \subset B^\alpha$ , (b)  $Q_K(p, q) = B^\alpha$  if and only if

$$\int_0^1 (1 - r^2)^{-2} K(-\log r) r dr < \infty. \quad (2)$$

Moreover,  $\|f\|_{B^\alpha} \lesssim \|f\|_{Q_K(p,q)}$ .

**Lemma 2.** Let  $0 < p < \infty$  and  $-2 < q < \infty$ ,  $K: [0, \infty) \rightarrow [0, \infty)$  is a non-decreasing continuous function such that (1) holds and  $f \in Q_K(p, q)$ .

Then

$$|f(z)| \lesssim \begin{cases} \|f\|_{Q_K(p,q)} & \text{if } 2 + q < p; \\ \log_2 \frac{2}{1 - |z|^2} \|f\|_{Q_K(p,q)} & \text{if } 2 + q = p; \\ \frac{\|f\|_{Q_K(p,q)}}{(1 - |z|^2)^{\frac{2+q}{p}-1}} & \text{if } 2 + q > p; \end{cases}$$

And

$$|f^{(n)}(z)| \lesssim \frac{\|f\|_{Q_K(p,q)}}{(1 - |z|^2)^{\frac{2+q}{p}+n-1}} \quad \text{if } n \geq 1.$$

The next lemma is an extension of a well known result on compactness of composition operators of the Hardy spaces  $H^p$  (see [1], proposition 3.11). The proof follows using the standard arguments of proposition 3.11 in [1]. We omit the details.

**Lemma 3.** Let  $0 < p < \infty$  and  $-2 < q < \infty$ ,  $K: [0, \infty) \rightarrow [0, \infty)$  is a non-decreasing continuous function such that (1) holds,  $\omega$  a radial weight,  $\psi \in H(\mathbb{D})$ ,  $n \in \mathbb{N}$  or  $n = 0$  and  $\varphi$  be a holomorphic self-map on  $\mathbb{D}$ . Then  $D_{\varphi,\psi}^n: Q_K(p, q) \rightarrow B_\omega$  is compact if and only if for every bounded sequence  $\{f_k\}$  in  $Q_K(p, q)$  which converges to zero on compact subset of  $\mathbb{D}$  as  $k \rightarrow \infty$ , we have  $\|D_{\varphi,\psi}^n f_k\|_{B_\omega} \rightarrow 0$  as  $k \rightarrow \infty$ .

The next two lemmas are inspired from lemmas 3.6 and 3.7 in [5].

**Lemma 4.** Let  $0 < p < \infty$  and  $-2 < q < \infty$  such that  $2 + q < p$  and  $K: [0, \infty) \rightarrow [0, \infty)$  be a non-decreasing continuous function such that (1) holds. Then every bounded sequence in  $Q_K(p, q)$  has a subsequence that converges uniformly on  $\overline{\mathbb{D}}$ .

Proof. Let  $\{f_m\}$  be a sequence in  $Q_K(p, q)$  and  $M$  be a positive number such that  $\|f_m\|_{Q_K(p,q)} \leq M$  for all  $m \in \mathbb{N}$ . Since  $Q_K(p, q) \subset \mathcal{B}^{(2+q)/p}$  and  $(2 + q)/p < 1$ , we have  $|f_m(z) - f_m(w)| \lesssim |z - w|^{(p-2-q)/p}$  for all  $z, w \in \mathbb{D}$  and for every  $m \in \mathbb{N}$ . Thus the family  $\{f_m: m \in \mathbb{N}\}$  is equicontinuous. Since  $\|f\|_\infty \lesssim \|f\|_{\mathcal{B}^{(2+q)/p}} \lesssim \|f\|_{Q_K(p,q)}$  for every analytic  $f$  on  $\mathbb{D}$ , the family  $\{f_m: m \in \mathbb{N}\}$  is bounded in  $C(\overline{\mathbb{D}})$ . Hence by Arzela-Ascoli Theorem the result follows.

**Lemma 5.** Let  $0 < p < \infty$  and  $-2 < q < \infty$  such that  $2 + q < p$ ,  $K: [0, \infty) \rightarrow [0, \infty)$  be a non-decreasing continuous function such that (1) holds and  $T: Q_K(p, q) \rightarrow X$  is a bounded linear operator, where  $X$  is a normed linear space. Then  $T: Q_K(p, q) \rightarrow X$  is compact if and only if  $\|Tf_m\| \rightarrow 0$  for every bounded sequence  $\{f_m\}$  in  $Q_K(p, q)$  that converges uniformly on  $\overline{\mathbb{D}}$ .

Proof. If  $T: Q_K(p, q) \rightarrow X$  is compact, then it is obvious that  $\|Tf_m\| \rightarrow 0$  for every bounded sequence  $\{f_m\}$  in  $Q_K(p, q)$  that converges uniformly on  $\overline{\mathbb{D}}$ . Conversely, suppose  $T: Q_K(p, q) \rightarrow X$  is not compact. Then there is a bounded sequence  $\{f_m\}$  in  $Q_K(p, q)$  such that  $\{Tf_m\}$  has no convergent subsequence. By Lemma 5,  $\{f_m\}$  has a subsequence  $\{g_m\}$  such that  $g_m \rightarrow g$  uniformly on  $\overline{\mathbb{D}}$ . Clearly,  $g \in Q_K(p, q)$  and sequence  $\{g_m - g\}$  is bounded in  $Q_K(p, q)$  and converges to zero uniformly on  $\overline{\mathbb{D}}$ . By

assumption  $\|T(g_m - g)\| \rightarrow 0$  as  $m \rightarrow \infty$ . Thus the subsequence  $\{T(g_m)\}$  of  $\{T(f_m)\}$  converges in  $X$ , a contradiction. Hence we are done.

**Theorem 1.** Let  $0 < p < \infty$  and  $-2 < q < \infty$ ,  $K: [0, \infty] \rightarrow [0, \infty]$  is a non-decreasing continuous function such that (1) holds,  $\omega$  a radial weight,  $\psi \in H(\mathbb{D})$ ,  $n \in \mathbb{N}$  or  $n = 0$  and  $\varphi$  be a holomorphic self-map on  $\mathbb{D}$ .

(1) If  $n \in \mathbb{N}$ , or  $n = 0$  and  $2 + q > p$ , then  $D_{\varphi, \psi}^n: Q_K(p, q) \rightarrow \mathcal{B}_\omega$  is compact if and only if  $\psi \in \mathcal{B}_\omega, \psi\varphi' \in \mathcal{A}_\omega$  and

$$(a) \lim_{|\varphi(z)| \rightarrow 1} \frac{\omega(z)|\psi'(z)|}{(1-|\varphi(z)|^2)^{\frac{2+q}{p} + n - 1}} = 0,$$

$$(b) \lim_{|\varphi(z)| \rightarrow 1} \frac{\omega(z)|\varphi'(z)\psi(z)|}{(1-|\varphi(z)|^2)^{\frac{2+q}{p} + n}} = 0.$$

(2) If  $2 + q < p$ , then  $D_{\varphi, \psi}^0: Q_K(p, q) \rightarrow \mathcal{B}_\omega$  is compact if and only if  $\psi \in \mathcal{B}_\omega, \psi\varphi' \in \mathcal{A}_\omega$  and

$$(c) \lim_{|\varphi(z)| \rightarrow 1} \frac{\omega(z)|\psi(z)\varphi'(z)|}{(1-|\varphi(z)|^2)^{\frac{2+q}{p}}} = 0.$$

(3) If  $2 + q = p$  and condition (2) of Lemma 1 holds, then  $D_{\varphi, \psi}^0: Q_K(p, q) \rightarrow \mathcal{B}_\omega$  is compact if and only if  $\psi \in \mathcal{B}_\omega, \psi\varphi' \in \mathcal{A}_\omega$  and

$$(d) \lim_{|\varphi(z)| \rightarrow 1} \frac{\omega(z)|\psi(z)\varphi'(z)|}{(1-|\varphi(z)|^2)^{\frac{2+q}{p}}} = 0,$$

$$(e) \lim_{|\varphi(z)| \rightarrow 1} \omega(z)|\psi'(z)| \log\left(\frac{2}{1-|\varphi(z)|^2}\right) = 0.$$

Proof. (1). Suppose that  $n \in \mathbb{N}$ , or  $n = 0$  and  $2 + q > p$  and  $D_{\varphi, \psi}^n: Q_K(p, q) \rightarrow \mathcal{B}_\omega$  is compact. Let  $\{z_k\}$  be a sequence in  $\mathbb{D}$  such that  $|\varphi(z_k)| \rightarrow 1$  as  $k \rightarrow \infty$ . Define

$$f_k(z) = \frac{1-|\varphi(z_k)|^2}{\left(1-\frac{\overline{\varphi(z_k)}z}{z}\right)^{\frac{2+q}{p}}} \quad (3)$$

And

$$g_k(z) = \frac{1-|\varphi(z_k)|^2}{\left(1-\frac{\overline{\varphi(z_k)}z}{z}\right)^{\frac{2+q}{p}}} - \frac{2+q}{2+q+np} \left(\frac{(1-|\varphi(z_k)|^2)^2}{(1-\varphi(z_k))^{\frac{2+q}{p}+1}}\right) \quad (4)$$

Then  $\{f_k\}$  and  $\{g_k\}$  are norm bounded sequences in  $Q_K(p, q)$  which converges to zero uniformly on compact subsets of  $\mathbb{D}$  as  $k \rightarrow \infty$ . By Lemma 3, we have

$$\lim_{k \rightarrow \infty} \|D_{\varphi, \psi}^n f_k\|_{\mathcal{B}_\omega} = 0 \quad \text{and} \quad \lim_{k \rightarrow \infty} \|D_{\varphi, \psi}^n g_k\|_{\mathcal{B}_\omega} = 0 \quad (5)$$

Thus we have that

$$\frac{p}{2+q+np} \prod_{j=1}^{n+1} \left(\frac{2+q}{p} + j - 1\right) \left|\overline{\varphi(z_k)}\right|^{n+1} \frac{\omega(z_k)|\psi(z_k)\varphi'(z_k)|}{|1-|\varphi(z_k)||^2 \frac{2+q}{p} + n} \|D_{\varphi, \psi}^n g_k\|_{\mathcal{B}_\omega} \quad (6)$$

And

$$\begin{aligned} \|D_{\varphi, \psi}^n f_k\|_{\mathcal{B}_\omega} \gtrsim & \prod_{j=1}^n \left(\frac{2+q}{p} + j - 1\right) \overline{\varphi(\lambda)}^n \frac{\omega(z_k)\psi'(\lambda)}{(1-|\varphi(\lambda)|^2)^{\frac{2+q}{p} + n - 1}} \\ & + \prod_{j=1}^{n+1} \left(\frac{2+q}{p} + j - 1\right) \overline{\varphi(\lambda)}^{n+1} \frac{\omega(z_k)\psi(\lambda)\varphi'(\lambda)}{(1-|\varphi(\lambda)|^2)^{\frac{2+q}{p} + n}} \end{aligned} \quad (7)$$

Taking  $k \rightarrow \infty$  on both sides of (6) and (7) and employing (5), we get (a) and (b). Moreover, as in Theorem 1, we can prove that  $\psi \in \mathcal{B}_\omega$  and  $\psi\varphi' \in \mathcal{A}_\omega$ . Conversely, suppose that  $\psi \in \mathcal{B}_\omega, \psi\varphi' \in \mathcal{A}_\omega$  and (a) and (b) hold. Let  $\{f_k\}$  be a sequence in  $Q_K(p, q)$  such that  $f_k \rightarrow 0$  uniformly on compact subsets of  $\mathbb{D}$  and  $\|f_k\|_{Q_K(p, q)} \lesssim 1$ . Then  $f_k^{(n)}$  and  $f_k^{(n+1)}$  converges to zero uniformly on compact subsets of  $\mathbb{D}$ . Since (a) and (b) hold, for every  $\epsilon > 0$ , there exist  $\delta \in (0, 1)$  such that

$$\frac{\omega(z)|\psi'(z)|}{(1 - |\varphi(z)|^2)^{\frac{2+q}{p}+n-1}} < \epsilon \text{ and } \frac{\omega(z)|\psi(z)\varphi'(z)|}{(1 - |\varphi(z)|^2)^{\frac{2+q}{p}+n}} < \epsilon,$$

when  $\delta < |\varphi(z)| < 1$ . Since  $\psi \in \mathcal{B}, \psi\varphi' \in \mathcal{A}_\infty^1$ , we have  $N_1 = \sup_{z \in \mathbb{D}} \omega(z)|\psi'(z)| < \infty$  and  $N_2 = \sup_{z \in \mathbb{D}} \omega(z)|\psi(z)\varphi'(z)| < \infty$ . This together with Lemma 3 yields

$$\begin{aligned} \|D_{\varphi,\psi}^n f_k\|_{\mathcal{B}_\omega} &\leq \sup_{z \in \mathbb{D}} \omega(z) |(D_{\varphi,\psi}^n f_k)'(z)| \\ &\leq \sup_{z \in \mathbb{D}} \omega(z) |\psi'(z) f_k^{(n)}(\varphi(z))| + \sup_{z \in \mathbb{D}} \omega(z) |\psi(z)\varphi'(z) f_k^{(n+1)}(\varphi(z))| \\ &\leq \sup_{|\varphi(z)| \leq \delta} \omega(z) |\psi'(z)| |f_k^{(n)}(\varphi(z))| + \sup_{|\varphi(z)| > \delta} \omega(z) |\psi'(z)| |f_k^{(n)}(\varphi(z))| \\ &\quad + \sup_{|\varphi(z)| \leq \delta} \omega(z) |\psi(z)\varphi'(z)| |f_k^{(n+1)}(\varphi(z))| \\ &\quad + \sup_{|\varphi(z)| > \delta} \omega(z) |\psi(z)\varphi'(z)| |f_k^{(n+1)}(\varphi(z))| \\ &\leq N_1 \sup_{|\varphi(z)| \leq \delta} |f_k^{(n)}(\varphi(z))| + \sup_{|\varphi(z)| > \delta} \frac{\omega(z)|\psi'(z)|}{(1 - |\varphi(z)|^2)^{\frac{2+q}{p}+n-1}} \\ &\quad + N_2 \sup_{|\varphi(z)| \leq \delta} |f_k^{(n+1)}(\varphi(z))| + \sup_{|\varphi(z)| > \delta} \frac{\omega(z)|\psi(z)\varphi'(z)|}{(1 - |\varphi(z)|^2)^{\frac{2+q}{p}+n-1}} \\ &\leq N_1 \sup_{|\varphi(z)| \leq \delta} |f_k^{(n)}(\varphi(z))| + N_2 \sup_{|\varphi(z)| \leq \delta} |f_k^{(n+1)}(\varphi(z))| + 2\epsilon \end{aligned}$$

From the arbitrariness of  $\epsilon$ , and the fact that  $f_k^{(n)}$  and  $f_k^{(n+1)}$  converges to zero uniformly on compact subsets of  $\mathbb{D}$ , we have  $\lim_{k \rightarrow \infty} \|D_{\varphi,\psi}^n f_k\|_{\mathcal{B}_\omega} = 0$  and so by Lemma 4,  $D_{\varphi,\psi}^n: Q_K(p, q) \rightarrow \mathcal{B}_\omega$  is compact.

(2). The necessity in condition (2) can be proved in the same way as the proof above. We omit the details. Assume that  $2 + q < p, \psi \in \mathcal{B}_\omega, \psi\varphi' \in \mathcal{A}_\omega$  and (c) holds. Let  $\{f_m\}$  be a bounded sequence in  $Q_K(p, q)$  which converges to zero uniformly on compact subsets of  $\mathbb{D}$ . To show that  $D_{\varphi,\psi}^0: Q_K(p, q) \rightarrow \mathcal{B}_\omega$  is compact, we need to show that if  $M = \sup_m \|f_m\|_{Q_K(p,q)} < \infty$  and  $f_m \rightarrow 0$  uniformly on compact subsets of  $\mathbb{D}$ , then  $\|D_{\varphi,\psi}^0 f_m\|_{\mathcal{B}_\omega} \rightarrow 0$  as  $m \rightarrow \infty$ . This amounts to show that

$$\sup_{z \in \mathbb{D}} \omega(z) |\psi(z)\varphi'(z) f_m'(\varphi(z))| \rightarrow 0 \tag{8}$$

And

$$\sup_{z \in \mathbb{D}} \omega(z) |\psi'(z) f_m(\varphi(z))| \rightarrow 0 \tag{9}$$

as  $n \rightarrow \infty$ . By Lemma 6, we have  $\sup_{z \in \bar{D}} |f_m(z)| \rightarrow 0$  as  $n \rightarrow \infty$ . So

$$\sup_{z \in \mathbb{D}} \omega(z) |\psi'(z) f_m(\varphi(z))| \leq \|\psi\|_{\mathcal{B}} \sup_{z \in \mathbb{D}} |f_m(z)| \rightarrow 0$$

as  $n \rightarrow \infty$ . Thus (33) holds. If  $|\varphi(z)| \leq r < 1$ , then

$$\omega(z) |\psi(z)\varphi'(z) f_m'(\varphi(z))| \leq \|\psi\varphi'\|_{\mathcal{A}_\omega} \max_{|z| \leq r} |f_m'(z)|$$

If  $|\varphi(z)| > r$ , then

$$\omega(z) |\psi(z)\varphi'(z) f_m'(\varphi(z))| \leq \frac{\omega(z) |\psi(z)\varphi'(z)|}{(1 - |\varphi(z)|^2)^{\frac{2+q}{p}}}$$

Thus

$$\begin{aligned} \sup_{z \in \mathbb{D}} \omega(z) |\psi(z)\varphi'(z) f_m'(\varphi(z))| &\leq \|\psi\varphi'\|_{\mathcal{A}_\omega} \max_{|z| \leq r} |f_m'(z)| \\ &\quad + \sup_{|\varphi(z)| > r} \frac{\omega(z) |\psi(z)\varphi'(z)|}{(1 - |\varphi(z)|^2)^{\frac{2+q}{p}}}. \end{aligned} \tag{10}$$

First letting  $n \rightarrow \infty$  and subsequently  $r$  increases to 1, in (10) obtains that (8) holds. (3). To prove (3), we assume that  $\{z_m\}$  is a sequence in  $\mathbb{D}$  for which  $|\varphi(z_m)| \rightarrow 1$  and consider the function

$$g_m(z) = \frac{3}{\mu_m} \left( \log \frac{2}{1 - \varphi(z_m)z} \right)^3 - \frac{2}{\mu_m^2} \left( \log \frac{2}{1 - \varphi(z_m)z} \right)^2$$

where  $\mu_m = \frac{\log \frac{2}{1 - |\varphi(z_m)|^2}}{1 - |\varphi(z_m)|^2}$ . Then

$$g'_m(z) = \frac{2\overline{\varphi(z_m)}}{1 - \overline{\varphi(z_m)}z} \left\{ \frac{1}{\mu_m} \left( \log \frac{2}{1 - \varphi(z_m)z} \right) - \frac{1}{\mu_m^2} \left( \log \frac{2}{1 - \varphi(z_m)z} \right)^2 \right\}.$$

By a direct calculation, we have  $\sup_m \|g_m\|_{\mathcal{B}} \leq C < \infty$ . Since condition (2) of Lemma 1 holds, we have  $g_m \in Q_K(p, q)$  and  $\sup_m \|f_m\|_{Q_K(p, q)}$  which converges to zero uniformly on compact subsets of  $\mathbb{D}$ . Note that

$$g'_m(\varphi(z_m)) = 0 \text{ and } g_m(\varphi(z_m)) = \log \left( \frac{2}{1 - |\varphi(z_m)|} \right).$$

So if  $D_{\varphi, \psi}^0: Q_K(p, q) \rightarrow \mathcal{B}_\omega$  is compact, then  $\|D_{\varphi, \psi}^0 g_m\|_{\mathcal{B}_\omega} \rightarrow 0$ , and consequently,

$$\omega(z_m) |\psi'(z_m)| \log \left( \frac{2}{1 - |\varphi(z_m)|^2} \right) = \omega(z_m) |(D_{\varphi, \psi}^n g_m)'(z_m)| \rightarrow 0$$

as  $n \rightarrow \infty$ . Conversely, suppose that  $\{f_m\}$  is a bounded sequence in  $Q_K(p, q)$  which converges to 0 uniformly on compact subsets of  $\mathbb{D}$ . Let  $M = \sup_m \|f_m(z)\|_{Q_K(p, q)} < \infty$ . By Lemma 2, we have

$$\sup_m |f_m(z)| \leq CM \log \frac{2}{1 - |z|^2}.$$

If  $|\varphi(z)| \leq r < 1$ , then

$$\omega(z) |\psi'(z) f_m(\varphi(z))| \leq \|\psi\|_{\mathcal{B}_\omega} \max_{|z| \leq r} |f_m(z)|$$

If  $|\varphi(z)| > r$ , then

$$\omega(z) |\psi'(z) f_m(\varphi(z))| \leq CM \omega(z) |\psi'(z)| \log \frac{2}{1 - |\varphi(z)|^2}$$

Thus

$$\sup_{z \in \mathbb{D}} \omega(z) |\psi'(z) f_m(z)| \leq \|\psi\|_{\mathcal{B}_\omega} \max_{|z| \leq r} |f_m(z)| + CM \omega(z) |\psi'(z)| \log \frac{2}{1 - |\varphi(z)|^2} \rightarrow 0$$

as  $n \rightarrow \infty$ . Similarly, we can prove that

$$\sup_{z \in \mathbb{D}} \omega(z) |\psi(z) \varphi'(z) f'_m(z)| \rightarrow 0$$

as  $n \rightarrow \infty$ . Hence  $\|D_{\varphi, \psi}^0 f_m\|_{\mathcal{B}, \omega} \rightarrow 0$  and this completes the proof

## References

1. Cowen CC, MacCluer BD. Composition operators on spaces of analytic functions, CRC Press Boca Raton, New York, 1995.
2. Hibscheiler RA, Portnoy N. Composition followed by differentiation between Bergman and Hardy spaces, Rocky Mountain J Math 2005;35:843-855.
3. Kotilainen M. On composition operators in  $Q_K$  type spaces, Journal of Function Spaces and Applications 2007;5:103-122.
4. Madigan K, Matheson A. Compact composition operators on the Bloch space, Trans. Amer. Math. Soc 1995;347:2679-2687.
5. Ohno S, Stroethoff K, Zhao R. Weighted composition operators between Bloch-type spaces, Rocky Mountain J Math 2003;33:191-215.
6. Shapiro JH. Composition operators and classical function theory, SpringerVerlag, New York 1993.
7. Sharma AK. Volterra composition operators between Bergman-Nevanlinna and Bloch-type spaces, Demonstratio Math 2009;42:607-618.

8. Sharma AK. Products of multiplication, composition and differentiation between weighted Bergman-Nevanlinna and Bloch-type spaces, Turk. J. Math 2011;35:275-291.
9. Sharma AK. Generalized composition operators on the Bergman space, Demonstratio Math 2011;44:359-372.
10. Sharma A, Sharma AK. Carleson measures and a class of generalized integration operators on the Bergman space, Rocky Mountain J. Math. (To appear).
11. Stević S, Sharma AK. Weighted composition operators between growth spaces of the upper-half plane, Util. Math 2011;84:265-272.
12. Stević S, Sharma AK. Weighted composition operators between Hardy and growth spaces of the upper half-plane, Appl. Math. Comput 2011;217:4928-4934.
13. Stević S, Sharma AK. Essential norm of composition operators between weighted Hardy spaces, Appl. Math. Comput 2011;217:6192-6197.
14. Stević S, Sharma AK, Sharma SD. Weighted composition operators from weighted Bergman spaces to weighted type spaces of the upper half-plane, Abstr. Appl. Anal 2011, 10, Article ID 989625.
15. Ueki SI. Weighted composition operators on some function spaces of entire functions, Bull. Belg. Math. Soc. Simon Stevin, Acta Sci. Math. (Szeged) 2010;17:343-353.
16. Yang W. Products of Composition and Differentiation Operators from  $Q_K(p, q)$  Spaces to Bloch-Type Spaces, Abstr. Appl. Anal 2009, 14, Article ID 741920.
17. Wulan H, Zhou J.  $Q_K$  type spaces of analytic functions, Journal of Function Spaces and Applications 2006;4:73-84.
18. Zhao R. On a general family of function space, Ann, Acad. Sci. Fenn. Math. Dissertationes 1996;105:1-56.
19. Zhu K. Bloch type spaces of analytic functions, Rocky Mountain J. Math 1993;23:1143-1177.